

**Clinical Quality Measure (CQM) Aggregate Data for EPs using the 2014 CQM Definitions**

State/ Territory/ District	VT								
Report As Of Date	03/31/16								
Total Unduplicated Providers(EPs) to ever receive payment for 2014 CQM definitions	324								
<b>INSTRUCTIONS:</b> Provide the statistical data listed in the headings below for the Aggregate Measure data for each Clinical Quality Measure selected by a provider during attestation. The statistical data Average and Standard Deviation is representative of the aggregate measure responses to meet the threshold. Please note that Exclusion or Exception count and percentage represents the providers who entered data for an exclusion or exception on the measure (when applicable) that was greater than 0. The # of unduplicated providers who selected column refers to the count of unique providers who selected the measure.									
Clinical Quality Measures	Description	Average	Standard Deviation	# of Exclusions	Exclusion %	# of Exceptions	Exception %	# of unduplicated providers who selected	# of providers who entered 0 in the denominator
<b>CMS 2/NQF 0418 /ACO 18 GPRO PREV-12</b> Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for clinical depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen.	7.5%	15.1%	8	8.8%	0	0.0%	91	0
<b>CMS 22/ACO 21 /GPRO PREV-11</b> Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented	Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated.	36.6%	17.8%	9	13.2%	0	0.0%	68	3
<b>CMS 68/NQF 0419 /ACO 39 /GPRO CARE-3</b> Documentation of Current Medications in the Medical Record	Percentage of visits for patients aged 18 years and older for which the eligible professional attests to documenting a list of current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration.	80.5%	28.4%			0	0.0%	280	9
<b>CMS 69/NQF 0421 /ACO 16 /GPRO PREV-9</b> Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up - Population 1	Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous six months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous six months of the current encounter.	30.6%	27.2%	4	2.2%			178	51
<b>CMS 122/NQF 0059 /ACO 27 /GPRO DM-2</b> Diabetes: Hemoglobin A1c Poor Control	Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period.	47.4%	42.6%	2	1.5%			131	19
<b>CMS 127/NQF 0043 /ACO 15 /GPRO PREV-8</b> Pneumonia Vaccination Status for Older Adults	Percentage of patients 65 years of age and older who have ever received a pneumococcal vaccine.	54.1%	30.0%					102	5
<b>CMS 130/NQF 0034 /ACO 19 /GPRO PREV-6</b> Colorectal Cancer Screening	Percentage of adults 50-75 years of age who had appropriate screening for colorectal cancer.	2.3%	11.9%	14	18.9%			74	9
<b>CMS 131/NQF 0055 /ACO 41 /GPRO DM-7</b> Diabetes: Eye Exam	Percentage of patients 18-75 years of age with diabetes who had retinal or dilated eye exam by an eye care professional during the measurement period or a negative retinal exam (no evidence of retinopathy) in the last 12 months prior to the measurement period.	16.6%	35.1%	0	0.0%			11	1
<b>CMS 138/NQF 0028 /ACO 17 /GPRO PREV-10</b> Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user.	76.6%	52.2%			0	0.0%	282	16
<b>CMS 139/NQF 0101 /ACO 13 /GPRO CARE-2</b> Falls: Screening for Future Fall Risk	Percentage of patients 65 years of age and older who were screened for future fall risk during the measurement period.	30.8%	43.3%			0	0.0%	14	5
<b>CMS 144/NQF 0083 /ACO 31 /GPRO HF-6</b> Heart Failure (HF): Beta-Blocker Therapy for Left Ventricular Systolic Dysfunction (LVSD)	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) <40% who were prescribed beta-blocker therapy either within a 12 month period when seen in the outpatient setting OR at each hospital discharge.	0.0%	0.0%			0	0.0%	0	0
<b>CMS 147/NQF 0041 /ACO 14 /GPRO PREV-7</b> Preventive Care and Screening: Influenza Immunization	Percentage of patients aged 6 months and older seen for a visit between October 1 and March 31 who received an influenza immunization OR who reported previous receipt of an influenza immunization.	30.3%	23.9%			0	0.0%	229	4
<b>CMS 159/NQF 0710 /ACO 40 /GPRO MH-1</b> Depression Remission at Twelve Months	Adult patients age 18 and older with major depression or dysthymia and an initial PHQ-9 score > 9 who demonstrate remission at twelve months defined as PHQ-9 score less than 5. This measure applies to both patients with newly diagnosis and existing depression who current PHQ-9 score indicates a need for treatment.	0.0%	0.0%	0	0.0%			3	3
<b>CMS 164/NQF 0068 /ACO 30 /GPRO IVD-2</b> Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antithrombotic	Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period, and who had documentation of use of aspirin or another antithrombotic during the measurement period.	36.8%	33.4%					27	5
<b>CMS 165/NQF 0018 /ACO 28 /GPRO HTN-2</b> Controlling High Blood Pressure	Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90 mmHg) during the measurement period.	54.4%	34.3%	20	14.7%			136	18